

The complete HIP brief is available from:

<https://www.fphighimpactpractices.org/briefs/postaborton-family-planning>

### References

1. World Health Organization, Department of Reproductive Health and Research (WHO/RHR); Johns Hopkins Bloomberg School of Public Health/Center for Communication Programs (CCP), Knowledge for Health Project. *Family Planning: A Global Handbook for Providers*. 2018 update. Baltimore and Geneva: CCP and WHO; 2018. <https://www.fphandbook.org/>. Accessed February 27, 2019.
2. Kangatharan C, Labram S, Bhattacharya S. Interpregnancy interval following miscarriage and adverse pregnancy outcomes: systematic review and meta-analysis. *Hum Reprod Update*. 2017;23(2):221-231. <https://doi.org/10.1093/humupd/dmw043>
3. International Federation of Gynecology and Obstetrics (FIGO); International Confederation of Midwives (ICM); International Council of Nurses (ICN); United States Agency for International Development (USAID); White Ribbon Alliance (WRA); Department for International Development (DFID); Bill & Melinda Gates Foundation. Post abortion family planning: A key component of post abortion care. Washington, DC: 2013. [https://www.figo.org/sites/default/files/uploads/project-publications/PAC-FP-Joint-Statement-November2013-final\\_printquality.pdf](https://www.figo.org/sites/default/files/uploads/project-publications/PAC-FP-Joint-Statement-November2013-final_printquality.pdf). Accessed February 27, 2019.
4. Huber D, Curtis C, Irani L, Pappa S, Arrington L. Postabortion care: 20 years of strong evidence on emergency treatment, family planning, and other programming components. *Glob Health Sci Pract*. 2016;4(3):481-494. <https://doi.org/10.9745/GHSP-D-16-00052>
5. Hossain A, Maddow-Zimet I, Ingerick M, Bhuiyan HU, Vlassoff M, Singh S. Access to and quality of menstrual regulation and postabortion care in Bangladesh: evidence from a survey of health facilities, 2014. New York: Guttmacher Institute; 2017. [https://www.guttmacher.org/sites/default/files/report\\_pdf/menstrual-regulation-postabortion-care-bangladesh\\_0.pdf](https://www.guttmacher.org/sites/default/files/report_pdf/menstrual-regulation-postabortion-care-bangladesh_0.pdf). Accessed February 27, 2019.
6. Pestvenidze E, Berdzuli N, Lomia N, Gagua T, Umikashvili L, Stray-Pedersen B. Repeat induced abortions in Georgia, characteristics of women with multiple pregnancy terminations: secondary analysis of the Reproductive Health Survey 2010. *Eur J Obstet Gynecol Reprod Biol*. 2016;205:85-90. <https://doi.org/10.1016/j.ejogrb.2016.08.020>
7. EngenderHealth. Post Abortion Care-Family Planning Annual Report for the Period October 1, 2016 - September 30, 2017. New York: EngenderHealth; 2017.
8. Zaidi S, Yasmin H, Hassan L, Khakwani M, Sami S, Abbas T. Replacement of dilation and curettage/evacuation by manual vacuum aspiration and medical abortion, and the introduction of postabortion contraception in Pakistan. *Int J Gynaecol Obstet*. 2014;126(suppl 1):S40-S44. <https://doi.org/10.1016/j.ijgo.2014.03.016>

9. Mutua MM, Achia TNO, Maina BW, Izugbara CO. A cross-sectional analysis of Kenyan postabortion care services using a nationally representative sample. *Int J Gynaecol Obstet.* 2017;138(3):276-282. <https://doi.org/10.1002/ijgo.12239>
10. Rocca CH, Puri M, Harper CC, Blum M, Dulal B, Henderson JT. Postabortion contraception a decade after legalization of abortion in Nepal. *Int J Gynaecol Obstet.* 2014;126(2):170-174. <https://doi.org/10.1016/j.ijgo.2014.02.020>
11. Borges AL, Monteiro RL, Hoga LA, Fujimori E, Chofakian CB, dos Santos OA. Post-abortion contraception: care and practices. *Rev Lat Am Enfermagem.* 2014;22(2):293-300. <https://dx.doi.org/10.1590%2F0104-1169.3240.2415>
12. High Impact Practices in Family Planning (HIP). Family planning high impact practices list. Washington, DC: USAID; 2018. <https://www.fphighimpactpractices.org/high-impact-practices-in-family-planning-list/>. Accessed February 27, 2019.
13. Sedgh G, Bearak J, Singh S, et al. Abortion incidence between 1990 and 2014: global, regional, and subregional levels and trends. *Lancet.* 2016; 388(10041):258-267. [https://doi.org/10.1016/S0140-6736\(16\)30380-4](https://doi.org/10.1016/S0140-6736(16)30380-4)
14. Sedgh G, Singh S, Shah IH, Ahman E, Henshaw S, Bankole A. Induced abortion: incidence and trends worldwide from 1995 to 2008. *Lancet.* 2012;379(9816):625-632. [https://doi.org/10.1016/S0140-6736\(11\)61786-8](https://doi.org/10.1016/S0140-6736(11)61786-8)
15. Singh S, Remez L, Sedgh G, Kwok L, Onda T. Abortion worldwide 2017: uneven progress and unequal access. New York: Guttmacher Institute; 2018. [https://www.guttmacher.org/sites/default/files/report\\_pdf/abortion-worldwide-2017.pdf](https://www.guttmacher.org/sites/default/files/report_pdf/abortion-worldwide-2017.pdf). Accessed February 27, 2019.
16. Lule E, Singh S, Chowdhury SA. Fertility regulation behaviors and their costs: contraception and unintended pregnancies in Africa and Eastern Europe and Central Asia. Washington, DC: World Bank; 2007. <http://siteresources.worldbank.org/HEALTHNUTRITIONANDPOPULATION/Resources/281627-1095698140167/FertilityRegulationsFinal.pdf>. Accessed February 27, 2019.
17. Jones RK, Singh S, Finer LB, Frohwirth LF. Repeat abortion in the United States. New York: Guttmacher Institute; 2006. Occasional Report 29. <http://www.guttmacher.org/pubs/2006/11/21/or29.pdf>. Accessed February 27, 2019.
18. Peipert JF, Madden T, Allsworth JE, Secura, GM. Preventing unintended pregnancies by providing no-cost contraception. *Obstet Gynecol.* 2012;120(6):1291-1297. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4000282/>
19. Kidder E, Sonneveldt E, Hardee K. Who receives PAC services? Evidence from 14 countries. Washington, DC: The Futures Group; 2004.
20. Vlassoff M, Singh S, Onda T. The cost of postabortion care in developing countries: a comparative analysis of four studies. *Health Policy Plan.* 2016; 31(8):1020-1030. <http://doi.org/10.1093/heapol/czw032>
21. Schreiber CA, Sober S, Ratcliffe S, Creinin MD. Ovulation resumption after medical abortion with mifepristone and misoprostol. *Contraception.* 2011;84(3):230-233. <https://doi.org/10.1016/j.contraception.2011.01.013>
22. Mahmoud GA, Byomy SS. Fertility awareness and family planning use among post abortion women in Egypt. *Life Sci J.* 2013;10(1):143-150. <https://www.doi.org/10.7537/marslsj100113.21>

23. David PH, Reichenbach L, Savelieva I, Vartapetova N, Potemkina R. Women's reproductive health needs in Russia: what can we learn from an intervention to improve post-abortion care? *Health Policy Plan.* 2007;22(2):83-94. <https://doi.org/10.1093/heapol/czm003>
24. Ganatra B, Gerdts C, Rossier C, et al. Global, regional, and subregional classification of abortions by safety, 2010-14: estimates from a Bayesian hierarchical model. *Lancet.* 2017;390(10110):2372-2381. [https://doi.org/10.1016/S0140-6736\(17\)31794-4](https://doi.org/10.1016/S0140-6736(17)31794-4)
25. Say L, Chou D, Gemmill A, et al. Global causes of maternal death: a WHO systematic analysis. *Lancet Glob Health.* 2014;2(6):e323-33. [http://dx.doi.org/10.1016/S2214-109X\(14\)70227-X](http://dx.doi.org/10.1016/S2214-109X(14)70227-X)
26. Tripney J, Kwan I, Bird, KS. Postabortion family planning counseling and services for women in low income countries: a systematic review. *Contraception.* 2013;87(1):17-25. <https://doi.org/10.1016/j.contraception.2012.07.014>
27. Centre de Formation et de Recherche en Sante de la Reproduction (CEFOREP). Introduction des soins obstetricaux d'urgence et de la planification familiale pour les patients presentant des complications liees a un avortement incomplete. Dakar: CEFORP; 1998.
28. Senlet P, Cagatay L, Ergin J, Mathis J. Bridging the gap: integrating family planning with abortion services in Turkey. *Int Fam Plan Perspect.* 2001;27(2):90-95. [https://www.guttmacher.org/sites/default/files/article\\_files/2709001.pdf](https://www.guttmacher.org/sites/default/files/article_files/2709001.pdf). Accessed March 15, 2019.
29. Benson J, Andersen K, Brahmi D, et al. What contraception do women use after abortion? An analysis of 319,385 cases from eight countries. *Glob Public Health.* 2018;13(1):35-50. <https://doi.org/10.1080/17441692.2016.1174280>
30. Banerjee SK, Gulati S, Andersen KL, Acre V, Warvadekar J, Navin D. Associations between abortion services and acceptance of postabortion contraception in six Indian states. *Stud Fam Plann.* 2015;46(4):387-403. <https://doi.org/10.1111/j.1728-4465.2015.00039.x>
31. Chukwumalu K, Gallagher MC, Baunach S, Cannon A. Uptake of postabortion care services and acceptance of postabortion contraception in Puntland, Somalia. *Reprod Health Matters.* 2017;25(51):48-57. <https://doi.org/10.1080/09688080.2017.1402670>
32. Pfitzer A, Hyjazi Y, Arnold B, et al. Inclusion of long-acting reversible contraceptives as an option for postabortion care clients in Guinea: results of an observational study. *Glob Health Sci Pract.* 2019;7(suppl 2). In press
33. Johnson BR, Ndhlovu S, Farr SL, Chipato T. Reducing unplanned pregnancy and abortion in Zimbabwe through postabortion contraception. *Stud Fam Plann.* 2002;33(2):195-202. <https://doi.org/10.1111/j.1728-4465.2002.00195.x>
34. Benson J, Huapaya V. Sustainability of postabortion care in Peru. New York: Population Council; 2002. 45 p. <https://www.popline.org/node/235774>. Accessed February 27, 2019.
35. Ottolenghi E, Riveros P, Blanding S. Assessment of the Bolivia Postabortion Care Community Mobilization Program. ACQUIRE Project; 2008. [http://www.acquireproject.org/archive/files/5.0\\_community\\_engagement\\_marketing\\_and\\_communications/5.2\\_resources/5.2.2\\_studies/cPAC\\_report\\_revised\\_final.pdf](http://www.acquireproject.org/archive/files/5.0_community_engagement_marketing_and_communications/5.2_resources/5.2.2_studies/cPAC_report_revised_final.pdf). Accessed on March 18, 2019.

36. Undie CC, Obare F, RamaRao S. Replication of the Community Mobilization for Postabortion Care (COMMPAC) model in Naivasha District, Rift Valley Province, Kenya: evaluation report. The RESPOND Project Study Series: Contributions to Global Knowledge Report No. 9. New York: EngenderHealth; 2012. [http://www.respond-project.org/pages/files/6\\_pubs/research-reports/Study9-COMMPAC-Final-Evaluation-December2012-Final-forweb.pdf](http://www.respond-project.org/pages/files/6_pubs/research-reports/Study9-COMMPAC-Final-Evaluation-December2012-Final-forweb.pdf). Accessed February 27, 2019.
37. Wickstrom J, Russell N, Escandon I. Engaging communities as partners in postabortion care: a desk review of the Community Postabortion Care Project in Nakuru, Kenya. ACQUIRE Evaluation and Research Studies No. 11. New York: The ACQUIRE Project/Engender Health; 2008. [http://www.acquireproject.org/archive/files/11.0\\_research\\_studies/er\\_study\\_11.pdf](http://www.acquireproject.org/archive/files/11.0_research_studies/er_study_11.pdf). Accessed February 27, 2019.
38. Abdel-Tawab N, Huntington D, Hassan EO, Youssef H, Nawar L. Effects of husband involvement in postabortion patients' recovery and use of contraception in Egypt. In: Huntington D, Piet-Pelon NJ, editors. *Postabortion Care: Lessons from Operations Research*. New York: Population Council; 1999. pp. 16-37.
39. Kiggundu C. Decentralizing integrated postabortion care in Uganda: a pilot training and support initiative for improving the quality and availability of integrated RH service. Kampala, Uganda: Ministry of Health, PRIME, Ipas, and DISH; 1999.
40. Population Council. Improving the counseling and medical care of post abortion patients in Egypt: final report. Cairo, Egypt: Population Council; 1995. p. 215.
41. Billings D, Victor A, Baird T, Taylor J, Ababio K, Ntow S. Midwives and comprehensive postabortion care in Ghana. In: Huntington D, Piet-Pelon N, eds. *Postabortion Care: Lessons from Operations Research*. New York: Population Council; 1999. pp. 141-158.
42. Extending Service Delivery (ESD). A description of the private nurse midwives networks (clusters) in Kenya: a best practice model. Best Practices Series Report No. 2. Washington, DC: Pathfinder International, Extending Service Delivery Project; 2007.
43. Basnet I, Clapham S, Shaka G, McCall M. Evolution of the postabortion care program in Nepal: the contribution of a national safe motherhood project. *Int J Gynaecol Obstet*. 2004;86(1):98-108. <https://doi.org/10.1016/j.ijgo.2004.03.014>
44. Wanjiru M, Askew I, Munguti N, RamaRao S, Homan R. Assessing the feasibility, acceptability and cost of introducing postabortion care in health centres and dispensaries in rural Tanzania. Washington, DC: Population Council; 2007. [http://pdf.usaid.gov/pdf\\_docs/PNADJ505.pdf](http://pdf.usaid.gov/pdf_docs/PNADJ505.pdf). Accessed on February 27, 2019.
45. Johnston HB, Gallo MF, Benson J. Reducing the costs to health systems of unsafe abortion: a comparison of four strategies. *J Fam Plann Reprod Health Care*. 2007;33(4):250-257. <http://dx.doi.org/10.1783/147118907782101751>
46. Brambila C et al. Estimating costs of postabortion services at Dr. Aurelio Valdivieso General Hospital, Oaxaca, Mexico. In: Huntington D, Piet-Pelon N, ed. *Postabortion Care: Lessons from Operations Research*. New York: Population Council; 1999. pp. 108-124.
47. Sultana F, Nahar Q, Marions L, Oliveras E. Effect of post-menstrual regulation family planning service quality on subsequent contraceptive use in Bangladesh. *Int J Gynaecol Obstet*. 2013;123(suppl 1):e38-e42. <http://dx.doi.org/10.1016/j.ijgo.2013.07.007>

48. Medina R, Vernon R, Mendoza I, Aguilar C. Expansion of postpartum/postabortion contraception in Honduras. New York: Population Council; 2001. [http://pdf.usaid.gov/pdf\\_docs/Pnacm374.pdf](http://pdf.usaid.gov/pdf_docs/Pnacm374.pdf). Accessed on February 27, 2019.
49. McDougall J, Feters T, Clark KA, Rathavy T. Determinants of contraceptive acceptance among Cambodian abortion patients. *Stud Fam Plann.* 2009;40(2): 123-32. <http://dx.doi.org/10.1111/j.1728-4465.2009.00195.x>
50. Savelieva I, Pile JM, Sacci I, Loganathan R. Postabortion family planning operations research study in Perm, Russia. New York: Population Council; 2003. [http://pdf.usaid.gov/pdf\\_docs/pnacu602.pdf](http://pdf.usaid.gov/pdf_docs/pnacu602.pdf). Accessed on February 27, 2019.
51. Gessesew A. Abortion and unwanted pregnancy in Adigrat Zonal Hospital, Tigray, North Ethiopia. *Afr J Reprod Health.* 2010;14(3):183-188. <http://www.bioline.org.br/pdf?rh10034>. Accessed on March 15, 2019.
52. Hakimi M, Hayati EN, Marlinawati VU, Winkvist A, Ellsberg MC. Silence for the sake of harmony: domestic violence and health in Central Java, Indonesia. Yogyakarta, Indonesia: Umea University; 2001. <https://www.popline.org/node/571345>. Accessed on February 27, 2019.
53. Heise L, Ellsberg M, Gottemoelle M. Ending violence against women. *Population Reports. Series L: Issues in World Health.* Baltimore, MD: Johns Hopkins School of Public Health, Population Information Program; 1999. <http://www.k4health.org/toolkits/info-publications/ending-violence-against-women>. Accessed on February 27, 2019.
54. Kaye DK, Mirembe FM, Bantebya G, Johansson A, Ekstrom AM. Domestic violence as risk factor for unwanted pregnancy and induced abortion in Mulago Hospital, Kampala, Uganda. *Trop Med Int Health.* 2006;11(1):90-101. <https://doi.org/10.1111/j.1365-3156.2005.01531.x>
55. Keeling J, Birch L, Green P. Pregnancy counseling clinic: a questionnaire survey of intimate partner abuse. *J Fam Plann Reprod Health Care.* 2004;30(3):165-168. <http://dx.doi.org/10.1783/1471189041261663>
56. Silverman JG, Gupta J, Decker MR, Kapur N, Raj A. Intimate partner violence and unwanted pregnancy, miscarriage, induced abortion, and stillbirth among a national sample of Bangladeshi women. *BJOG.* 2007;114(10):1246-1252. <https://doi.org/10.1111/j.1471-0528.2007.01481.x>
57. World Health Organization (WHO). Responding to intimate partner violence and sexual violence against women: WHO clinical and policy guidelines. Geneva: WHO; 2013. <https://www.who.int/reproductivehealth/publications/violence/9789241548595/en/>. Accessed on February 27, 2019.
58. Fikree FF, Mugore S, Forrester H. Strengthening postabortion family planning in Burkina Faso: pre-service and in-service training on postabortion care. Washington, DC: Pathfinder International, Evidence to Action for Strengthened Reproductive Health; 2014. <https://www.e2aproject.org/wp-content/uploads/strengthening-pac-fp-burkina.pdf>. Accessed on February 27, 2019.
59. Fikree FF, Mugore S, Forrester H. Strengthening postabortion family planning in Senegal: maintaining and enhancing postabortion care services. Washington, DC: Pathfinder International, Evidence to Action for Strengthened Reproductive Health; 2014. <https://www.e2aproject.org/wp-content/uploads/strengthening-pac-fp-senegal.pdf>. Accessed on February 27, 2019.

60. EngenderHealth. Health system and patient cost for scaling up post abortion and post abortion care family planning services in Tanzania, 2018. New York: EngenderHealth; [2018].
61. Evens E, Otieno-Masaba R, Eichleay M, et al. Post-abortion care services for youth and adult clients in Kenya: a comparison of services, client satisfaction and provider attitudes. *J Biosoc Sci.* 2014;46(1):1-15. <https://doi.org/10.1017/S0021932013000230>
62. Alemayehu T, Biru D, Hendrickson C, Brahmi D, Feters T. An evaluation of postabortion contraceptive uptake following a youth-friendly services intervention in Ethiopia. Chapel Hill, NC: Ipas; 2016. [https://endabortionstigma.org/-/media/Files/PubsManual/PMDETHE16.ashx?utm\\_source=resource&utm\\_medium=meta&utm\\_campaign=PMDETHE16](https://endabortionstigma.org/-/media/Files/PubsManual/PMDETHE16.ashx?utm_source=resource&utm_medium=meta&utm_campaign=PMDETHE16). Accessed on February 27, 2019.
63. Hainsworth G, Boyce C, Israel E. Assessment of youth-friendly postabortion care services: a global tool for assessing and improving postabortion care for youth. Watertown, MA: Pathfinder International; 2008. [http://www2.pathfinder.org/site/DocServer/YFPAC\\_assessment\\_tool.pdf?docID=13121](http://www2.pathfinder.org/site/DocServer/YFPAC_assessment_tool.pdf?docID=13121). Accessed on February 27, 2019.
64. Interagency Working Group on Reproductive Health in Crisis (IAWG). Inter-Agency Field Manual on Reproductive Health in Humanitarian Settings. IAWG; 2018. <http://iawg.net/iafm/>. Accessed March 15, 2019.
65. Vu A, Adam A, Wirtz A, et al. The prevalence of sexual violence among female refugees in complex humanitarian emergencies: a systematic review and meta-analysis. *PLoS Curr.* 2014;6:pii:ecurrents.dis.835f10778fd80ae031aac12d3b533ca7. <https://doi.org/10.1371/currents.dis.835f10778fd80ae031aac12d3b533ca7>
66. Nielsen KK, Lusiola G, Kananma J, Bantamby J, Kikumbih N, Rasch V. Expanding comprehensive postabortion care to primary health facilities in Geita District, Tanzania. *Afr J Reprod Health.* 2009;13(2):129-38. <http://www.bioline.org.br/pdf?rh09029>. Accessed March 15, 2019.
67. Rasch V, Massawe S, Yambesi F, Bergstrom S. Acceptance of contraceptives among women who had an unsafe abortion in Dar es Salaam. *Trop Med Int Health.* 2004;9(3):399-405. <https://doi.org/10.1111/j.1365-3156.2004.01197.x>