

Methods for literature search, information sources, abstraction and synthesis**Search Strategy**

To compile the list of postabortion family planning (PAFP) documents meeting inclusion criteria, a literature search was conducted using bibliographic databases and hand searching of online websites for journal and grey literature. The period of review focused on documents published after the previous HIP brief on PAFP was developed in 2012 through April 2018.

Electronic Databases

The following electronic databases were searched:

- PubMed
- Scopus
- POPLINE
- Cochrane Database of Systematic Reviews (CDSR)
- African Journals Online
- Cumulative Index of Nursing and Allied Health (CINAHL)
- African Journals Online
- Science Direct
- Inter-Science (Wiley)
- Web of Science


In addition, online searches of nineteen websites containing relevant PAFP integration interventions/information were conducted.

Search Terms

Electronic database searching was conducted based on combinations of postabortion, family planning, and geographical concepts. Additional limits for publication date, language, and availability of abstract were also used.

Concept	Terms	String
Postabortion	postabortion, post abortion, post-abortion	postabortion OR post*abortion OR "post abortion"
Family Planning	family planning, contraception, contraceptives	"family planning" OR contracept*
Location	developing country, developing nation, low income country, low-income country, emergent nation, transitional country, limited resource country, limited-resource country, (country list)	low income country* OR developing country* OR developing nation* OR emergent nation*, transitional country*, limited resource country*
Additional Limits	after 2013, English, no commentary, abstract available for screening	

Document Flow



Bibliography

Below is a complete list of citations identified from the literature search process described above. A subset of these references were used to support the development of the HIP Brief. The complete brief and reference listing is available from: <https://www.fphighimpactpractices.org/briefs/postabortion-family-planning/>

- Aimable MM. Assessing immediate postpartum and postabortion contraception KAP among healthcare professionals in Kigali. *Obstetrics and Gynecology*. 2017;129:18S-18S. https://journals.lww.com/greenjournal/Abstract/2017/05001/Assessing_Immediate_Postpartum_and_Postabortion.58.aspx
- Alemayehu T, Hendrickson C, Brahmi D, Desta D, Fetter T. An evaluation of postabortion contraceptive uptake following a youth-friendly service intervention in Ethiopia. *Contraception*. 2013;88(3):447-448. [http://www.contraceptionjournal.org/article/S0010-7824\(13\)00360-0/abstract](http://www.contraceptionjournal.org/article/S0010-7824(13)00360-0/abstract)
- Amo-Adjei J, Darteh EKM. Unmet/met need for contraception and self-reported abortion in Ghana. *Sex Reprod Healthc*. 2017;13:118-124. <https://dx.doi.org/10.1016/j.srhc.2017.02.002>
- Arambepola C, Rajapaksa LC, Galwaduge C. Usual hospital care versus post-abortion care for women with unsafe abortion: a case control study from Sri Lanka. *BMC Health Services Research*. 2014;14:470. <https://bmchealthservres.biomedcentral.com/track/pdf/10.1186/1472-6963-14-470?site=bmchealthservres.biomedcentral.com>

- Banerjee SK, Gulati S, Andersen KL, Acre V, Warvadekar J, Navin D. Associations between abortion services and acceptance of postabortion contraception in six Indian States. *Studies in Family Planning*. 2015;46(4):387-403. <https://onlinelibrary.wiley.com/doi/full/10.1111/j.1728-4465.2015.00039.x>
- Barot S. Implementing postabortion care programs in the developing world: Ongoing challenges. *Guttmacher Policy Review*. 2014;17(1):22-28. https://www.guttmacher.org/sites/default/files/article_files/gpr170122.pdf
- Begum F, Zaidi S, Fatima P, Shamsuddin L, Anowar-ul Azim AKM. Improving manual vacuum aspiration service delivery, introducing misoprostol for cases of incomplete abortion, and strengthening postabortion contraception in Bangladesh. *International Journal of Gynecology & Obstetrics*. 2014;126 (Suppl 1):S31-S35. <https://www.sciencedirect.com/science/article/pii/S0020729214001544?via%3Dihub>
- Benson J, Andersen K, Brahmi D, et al. What contraception do women use after abortion? An analysis of 319,385 cases from eight countries. *Global Public Health*. 2018;13(1):35-50. <https://www.tandfonline.com/doi/full/10.1080/17441692.2016.1174280>
- Biswas KK, Pearson E, Shahidullah SM, Sultana S, Chowdhury R, Andersen KL. Integrating postabortion care, menstrual regulation and family planning services in Bangladesh: a pre-post evaluation. *Reproductive Health*. 2017;14(1):37. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5346262/pdf/12978_2017_Article_298.pdf
- Blanco MN, Taquette SR, Monteiro DL, Miranda FR. Postnatal and postabortion care during adolescence in the National Health System in Rio de Janeiro, Brazil. *International Journal of Gynaecology & Obstetrics*. 2014;126(2):126-129. <http://europepmc.org/abstract/MED/24825499>
- Borges AL, Monteiro RL, Hoga LA, Fujimori E, Chofakian CB, dos Santos OA. Post-abortion contraception: Care and practices. *Revista Latino-Americana de Enfermagem*. 2014;22(2):293-300. <https://www.popline.org/node/658896>
- Borges AL, OlaOlorun F, Fujimori E, Hoga LA, Tsui AO. Contraceptive use following spontaneous and induced abortion and its association with family planning services in primary health care: Results from a Brazilian longitudinal study. *Reproductive Health*. 2015;12:94. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4606494/pdf/12978_2015_Article_87.pdf
- Campbell OMR, Aquino EML, Vwalika B, Gabrysch S. Signal functions for measuring the ability of health facilities to provide abortion services: An illustrative analysis using a health facility census in Zambia. *BMC Pregnancy Childbirth*. 2016;16:105. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4868015/pdf/12884_2016_Article_872.pdf
- Center for Health Market Innovations. K-MET Post Abortion Care Network. <https://healthmarketinnovations.org/program/k-met-post-abortion-care-network>
- Center for Health Market Innovations. Women's Health Program (Pehel). <https://healthmarketinnovations.org/program/womens-health-program-pehel>
- Che Y, Liu X, Zhang B, Cheng L. Oral contraception following abortion: A systematic review and meta-analysis. *Medicine*. 2016;95(27):e3825. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5058789/pdf/medi-95-e3825.pdf>
- Chukwumalu K, Gallagher MC, Baunach S, Cannon A. Uptake of postabortion care services and acceptance of postabortion contraception in Puntland, Somalia. *Reproductive Health Matters*. 2017;25(51):48-57. <https://www.tandfonline.com/doi/full/10.1080/09688080.2017.1402670>

- Demeke D, Alemayehu T, Gelaso M. Integration of abortion care and family planning services: Family planning acceptance, trend and method preference among abortion care clients in Ipas-supported facilities. *Contraception*. 2013;88(3):447-447. [http://www.contraceptionjournal.org/article/S0010-7824\(13\)00359-4/abstract](http://www.contraceptionjournal.org/article/S0010-7824(13)00359-4/abstract)
- DePineres T, Benitez J, Becker D, Gil L. Randomized-controlled trial of contraceptive use at 3-and 6-months post-abortion. *Contraception*. 2014;89(5):481. <https://doi.org/10.1016/j.contraception.2014.02.024>
- DePineres T, Benitez J, Gil L. Pilot-randomized controlled trial of contraceptive use at 3 months postabortion. *Contraception*. 2013;88:451. <http://cochranelibrary-wiley.com/o/cochrane/clcentral/articles/899/CN-01006899/frame.html>
- Desta D, Hendrickson C, Brahmi D, et al. Postabortion contraception method mix and uptake in Ipas-supported health facilities in Ethiopia. *Contraception*. 2014;90(3):316. [http://www.contraceptionjournal.org/article/S0010-7824\(14\)00422-3/abstract](http://www.contraceptionjournal.org/article/S0010-7824(14)00422-3/abstract)
- Esber A, Foraker RE, Hemed M, Norris A. Partner approval and intention to use contraception among Zanzibari women presenting for post-abortion care. *Contraception*. 2014;90(1):23-28. <https://doi.org/10.1016/j.contraception.2014.03.006>
- Evens E, Otieno-Masaba R, Eichleay M, et al. Post-abortion care services for youth and adult clients in Kenya: A comparison of services, client satisfaction and provider attitudes. *Journal of Biosocial Science*. 2014;46:1-15. <https://www.cambridge.org/core/journals/journal-of-biosocial-science/article/postabortion-care-services-for-youth-and-adult-clients-in-kenya-a-comparison-of-services-client-satisfaction-and-provider-attitudes/6AB3530E5A693872AD6D1552B5718E92>
- FIGO, ICM, ICN, USAID, WRA, DFID, Bill and Melinda Gates Foundation. Post abortion family planning: A key component of post abortion care.” November 2013. http://www.respond-project.org/pages/files/6_pubs/advocacy-materials/PAC-FP-Joint-Statement-November2013-final.pdf
- Fikree FF, Mugore S, Forrester H. Strengthening postabortion family planning in Burkina Faso: Pre-service and in-service training on postabortion care. Washington, DC: Pathfinder International, Evidence to Action for Strengthened Reproductive Health [E2A];2014. <https://www.e2aproject.org/publication/strengthening-postabortion-family-planning-in-burkina-faso/>
- Fikree FF, Mugore S, Forrester H. Strengthening postabortion family planning in Guinea: Sustained technical assistance Washington, DC: Pathfinder International, Evidence to Action for Strengthened Reproductive Health [E2A];2014. <https://www.e2aproject.org/publication/strengthening-postabortion-family-planning-in-guinea/>
- Fikree FF, Mugore S, Forrester H. Strengthening postabortion family planning in Senegal: Maintaining and enhancing postabortion care services. Washington, DC: Pathfinder International, Evidence to Action for Strengthened Reproductive Health [E2A];2014. <https://www.e2aproject.org/wp-content/uploads/strengthening-pac-fp-senegal.pdf>
- Fikree FF, Mugore S, Forrester H. Strengthening postabortion family planning in Togo: Mobilizing resources. Washington, DC: Pathfinder International, Evidence to Action for Strengthened Reproductive Health [E2A];2014. <https://www.e2aproject.org/wp-content/uploads/strengthening-pac-fp-togo.pdf>
- Fikree FF, Mugore S, Forrester HA. Postabortion care: Assessment of postabortion care services in four Francophone West Africa countries. Washington, DC: Pathfinder International, Evidence to Action for Strengthened Reproductive Health [E2A];2014. <https://www.e2aproject.org/publication/assessment-of-postabortion-care-in-four-francophone-west-africa-countries/>

- Gemzell-Danielsson K, Kopp Kallner H, Faundes A. Contraception following abortion and the treatment of incomplete abortion. International Journal of Gynaecology and Obstetrics. 2014;126(Suppl 1):S52-S55. <https://www.sciencedirect.com/science/article/pii/S0020729214001532?via%3Dihub>
- Gupta S, Prateek S, Sinha R, Shamsunder S, Mittal MK. IUCD inserted after first trimester abortion: An observational study of 100 cases. Journal of Obstetrics and Gynaecology. 2016;36(8):1067-1068. <https://doi.org/10.1080/01443615.2016.1196481>
- Hofmeyr GJ, Singata-Madlik M, Lawrie TA, Bergel E, Temmerman M. Effects of the copper intrauterine device versus injectable progestin contraception on pregnancy rates and method discontinuation among women attending termination of pregnancy services in South Africa: A pragmatic randomized controlled trial. Reproductive Health. 2016;13(1):42. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4835872/pdf/12978_2016_Article_153.pdf
- Hossain A, Maddow-Zimet, Ingerick M, Bhuiyan HU, Vlassoff M, Singh S. Access to and quality of menstrual regulation and postabortion care in Bangladesh: Evidence from a survey of health facilities, 2014. New York: Guttmacher Institute; 2017. https://www.guttmacher.org/sites/default/files/report_pdf/menstrual-regulation-postabortion-care-bangladesh_0.pdf
- Hou SP, Zhu WL, Li SM, Teng YC. Acceptance and continuation of contraceptive methods immediate postabortion. Gynecologic and Obstetric Investigation. 2017;82:86-95. <https://www.karger.com/Article/Abstract/445292>
- Huber D, Curtis C, Irani L, Pappa S, Arrington L. Postabortion care: 20 Years of strong evidence on emergency treatment, family planning, and other programming components. Global Health Science & Practice. 2016;4(3):481-494. <http://www.ghspjournal.org/content/4/3/481>
- Jiang H, Xu J, Richards E, et al. Opportunities, challenges and systems requirements for developing post-abortion family planning services: Perceptions of service stakeholders in China. PLoS One. 2017;12(10):e0186555. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5646849/pdf/pone.0186555.pdf>
- Kathpalia SK. Acceptance of family planning methods by induced abortion seekers: An observational study over five years. Medical Journal Armed Forces India. 2016;72(1):8-11. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4723710/pdf/main.pdf>
- Macha S, Muyuni M, Nkonde S, Faundes A. Increasing access to legal termination of pregnancy and postabortion contraception at the University Teaching Hospital, Lusaka, Zambia. International Journal of Gynaecology and Obstetrics. 2014;126(Suppl 1):S49-S51. <https://www.sciencedirect.com/science/article/pii/S002072921400160X>
- Macha S, Muyuni M, Nkonde S, Padua KS, Faundes A. Retrospective observational study of interventions to improve acceptance of long-acting reversible contraception following abortion at Lusaka Teaching Hospital, Zambia. International Journal of Gynecology and Obstetrics. 2018;141(1):32-36. <https://obgyn.onlinelibrary.wiley.com/doi/abs/10.1002/ijgo.12428>
- Maxwell L, Voetagbe G, Paul M, Mark A. Does the type of abortion provider influence contraceptive uptake after abortion? An analysis of longitudinal data from 64 health facilities in Ghana. BMC Public Health. 2015;15:586. <https://bmcpublichealth.biomedcentral.com/track/pdf/10.1186/s12889-015-1875-2?site=bmcpublichealth.biomedcentral.com>

- Mayi-Tsonga S, Obiang PA, Minkobame U, Ngouafé D, Ambounda N, De Souza MH. Introduction of postabortion contraception, prioritizing long-acting reversible contraceptives, in the principal maternity hospital of Gabon. International Journal of Gynecology and Obstetrics. 2014;126(Suppl 1):S45-S48. <https://www.sciencedirect.com/science/article/pii/S0020729214001623>
- Mugore S, Kassouta NT, Sebikali B, Lundstrom L, Saad A. Improving the quality of Postabortion Care Services in Togo Increased Uptake of Contraception. Global Health Science & Practice. 2016;4(3):495-505. <http://www.ghspjournal.org/content/4/3/495>
- Mutua MM, Achia TNO, Maina BW, Izugbara CO. A cross-sectional analysis of Kenyan postabortion care services using a nationally representative sample. International Journal of Gynecology and Obstetrics. 2017;138(3):276-282. <https://obgyn.onlinelibrary.wiley.com/doi/epdf/10.1002/ijgo.12239>
- Nozar F, Greif D, Coppola F, Fiol V, Briozzo L. Legal termination of pregnancy as an opportunity for expanding postabortion contraception: Experience at the Pereira Rossell Hospital, Montevideo, Uruguay. International Journal of Gynecology and Obstetrics. 2016;134(Suppl 1):S28-S30. <https://www.sciencedirect.com/science/article/pii/S0020729216302508>
- Okusanya BO, Oduwole O, Effa EE. Immediate postabortal insertion of intrauterine devices. Cochrane Database of Systematic Reviews. 2014;7:CD001777. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD001777.pub4/full>
- Onyegbule OA, Onumajuru CC, Ododo NA, Idih EE, Onyegbule PC. Uptake of post-abortion contraception among women who had manual vacuum aspiration at Federal Medical Centre, Owerri: A 5-year review. Tropical Journal of Obstetrics and Gynaecology. 2016;33(2):228-231. <http://www.tjogonline.com/article.asp?issn=0189-5117;year=2016;volume=33;issue=2;spage=228;epage=231;aulast=Onyegbule>
- Padmadas SS, Lyons-Amos M, Thapa S. Contraceptive behavior among women after abortion in Nepal. International Journal of Gynaecology and Obstetrics. 2014;127(2):132-137. <https://obgyn.onlinelibrary.wiley.com/doi/full/10.1016/j.ijgo.2014.05.012>
- Pathfinder International. Evidence to Action for Strengthened Reproductive Health [E2A]. E2A in Togo: improving the quality of postabortion care services. Washington, DC: E2A; 2015. <https://www.e2aproject.org/wp-content/uploads/e2a-in-togo-improving-pac.pdf>
- Paul M, Iyengar S, Essén B, et al. Does mode of follow-up influence contraceptive use after medical abortion in a low-resource setting? Secondary outcome analysis of a non-inferiority randomized controlled trial. BMC public health. 2016;16(1):1087. <https://bmcpublichealth.biomedcentral.com/track/pdf/10.1186/s12889-016-3726-1>
- Pearson E, Biswas KK, Andersen KL, et al. Correlates of contraceptive use 4 months postabortion: Findings from a prospective study in Bangladesh. Contraception. 2017;95(3):279-287. <https://www.sciencedirect.com/science/article/pii/S0010782415301797>
- Pearson E, Biswas KK, Chowdhury R, et al. Service delivery correlates of choosing short-acting contraceptives at the time of uterine evacuation in Bangladesh. International Perspectives on Sexual and Reproductive Health. 2017;43(2):67-74. <http://dx.doi.org/10.1363/43e3817>
- Pestvenidze E, Berdzuli N, Lomia N, Gagua T, Umikashvili L, Stray-Pedersen B. Repeat induced abortions in Georgia, characteristics of women with multiple pregnancy terminations: Secondary analysis of the Reproductive Health Survey 2010. European Journal of Obstetrics, Gynecology, and Reproductive Biology. 2016;205:85-90. [http://www.ejog.org/article/S0301-2115\(16\)30868-5/fulltext](http://www.ejog.org/article/S0301-2115(16)30868-5/fulltext)

- Pilato A, Betstadt S. PACS: Post-abortion contraception study. Contraception. 2014;90(2):199. [http://www.contraceptionjournal.org/article/S0010-7824\(14\)00316-3/fulltext](http://www.contraceptionjournal.org/article/S0010-7824(14)00316-3/fulltext)
- Prata N, Holston M, Fraser A, Melkamu Y. Contraceptive use among women seeking repeat abortion in Addis Ababa, Ethiopia. African Journal of Reproductive Health. 2013;17(4):56-65. <https://www.ajol.info/index.php/ajrh/article/viewFile/98374/87660>
- Puri M, Henderson JT, Harper CC, Blum M, Joshi D, Rocca CH. Contraceptive discontinuation and pregnancy post-abortion in Nepal: a longitudinal cohort study. Contraception. 2015;91(4):301-307. [http://www.contraceptionjournal.org/article/S0010-7824\(14\)00820-8/fulltext](http://www.contraceptionjournal.org/article/S0010-7824(14)00820-8/fulltext)
- Rees H, Pillay D, Mullick S, Chersich M. Strengthening implant provision and acceptance in South Africa with the 'Any woman, any place, any time' approach: An essential step towards reducing unintended pregnancies. Southern African Medical Journal. 2017;107(11):939-944. <http://www.samj.org.za/index.php/samj/article/view/12117>
- Rob U, Hena IS, Sultana N, Hossain MI, Yasmin R, Das TR, Ahmed FU. Reducing unsafe menstrual regulation through medication in Bangladesh. STEP UP Policy Brief. Dhaka, Bangladesh: Population Council; 2014. http://www.popcouncil.org/uploads/pdfs/2014STEPUP_BangladeshMRM_PolicyBrief.pdf
- Rocca CH, Puri M, Harper CC, Blum M, Dulal B, Henderson JT. Postabortion contraception a decade after legalization of abortion in Nepal. International Journal of Gynecology and Obstetrics. 2014;126(2):170-174. <https://obgyn.onlinelibrary.wiley.com/doi/abs/10.1016/j.ijgo.2014.02.020>
- Rogers C, Dantas JAR. Access to contraception and sexual and reproductive health information post-abortion: a systematic review of literature from low- and middle-income countries. Journal of Family Planning and Reproductive Health Care. 2017;43(4):309-318. <http://srh.bmj.com/content/43/4/309.long>
- Rominski SD, Morhe ES, Lori J. Post-abortion contraception choices of women in Ghana: A one-year review. Global Public Health. 2015;10(3):345-353. <http://aphrc.org/abortionresearch/index.php/component/k2/item/750-post-abortion-contraception-choices-of-women-in-ghana-a-one-year-review>
- Samuel M, Fetters T, Desta D. Strengthening postabortion family planning services in Ethiopia: Expanding contraceptive choice and improving access to long-acting reversible contraception. Global Health Science & Practice. 2016;4(Suppl 2):S60-72. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4990163/pdf/S60.pdf>
- Sathar Z, Rashida G, Shah Z, Singh S, Woog V. Unsafe abortion and postabortion care in Pakistan. Issues in Brief. 2013;(4):1-8. <https://www.guttmacher.org/sites/default/files/pdfs/pubs/FB-UAPCIP.pdf>
- Sebastian MP, Khan ME, Sebastian D. Unintended pregnancy and abortion in India: Country profile report with focus on Bihar, Madhya Pradesh and Odisha. New Delhi, India, Population Council, 2013. http://www.popcouncil.org/uploads/pdfs/2014STEPUP_IndiaCountryProfile.pdf
- Smith C, Ly S, Uk V, Warnock R, Edwards P, Free C. Process evaluation of a mobile phone-based intervention to support post-abortion contraception in Cambodia. Contraception and Reproductive Medicine. 2017;2:16. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5683466/pdf/40834_2017_Article_43.pdf
- Smith C, Ly S, Uk V, Warnock R, Free C. Women's views and experiences of a mobile phone-based intervention to support post-abortion contraception in Cambodia. Reproductive Health. 2017;14(1):72. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5469016/pdf/12978_2017_Article_329.pdf
- Smith C, Ngo TD, Gold J, et al. Effect of a mobile phone-based intervention on post-abortion contraception: a randomized controlled trial in Cambodia. Bull World Health Organ. 2015;93(12):842-850a. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4669734/pdf/BLT.15.160267.pdf>

- Smith C, Vannak U, Sokhey L, Ngo T, Gold J, Free C. Mobile technology for improved family planning (MOTIF): The development of a mobile phone-based (mHealth) intervention to support post-abortion family planning (PAFP) in Cambodia. *Reproductive Health*. 2016;13(1):1. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4700587/pdf/12978_2015_Article_112.pdf
- Spring H, Datta S, Sapkota S. Using behavioral science to design a peer comparison intervention for postabortion family planning in Nepal. *Frontiers in Public Health*. 2016;4:123. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4914549/pdf/fpubh-04-00123.pdf>
- Steinberg JR, Tschan JM, Henderson JT, Drey EA, Steinauer JE, Harper CC. Psychological distress and post-abortion contraceptive method effectiveness level chosen at an urban clinic. *Contraception*. 2013;88(6):717-724. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3844792/pdf/nihms-524298.pdf>
- Sultana F, Nahar Q, Marions L, Oliveras E. Effect of post-menstrual regulation family-planning service quality on subsequent contraceptive use in Bangladesh. *International Journal of Gynecology & Obstetrics*. 2013;123:e38-e42. <https://obgyn.onlinelibrary.wiley.com/doi/abs/10.1016/j.ijgo.2013.07.007>
- Tang L, Wu S, Li J, Wang K, Xu J, Temmerman M, Zhang WH. Post-abortion family planning counselling practice among abortion service providers in China: A nationwide cross-sectional study. *European Journal of Contraception and Reproductive Health Care*. 2017;22(1):24-29. <https://www.tandfonline.com/doi/abs/10.1080/13625187.2016.1255939?scroll=top&needAccess=true&journalCode=iejc20>
- Tesfaye G, Oljira L. Post abortion care quality status in health facilities of Guraghe zone, Ethiopia. *Reproductive Health*. 2013;10:35. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3726452/pdf/1742-4755-10-35.pdf>
- Tripney J, Kwan I, Bird KS,. Postabortion family planning counseling and services for women in low-income countries: a systematic review. *Contraception*. 2013;87(1):17-25. <https://www.ncbi.nlm.nih.gov/pubmed/22974595>
- Undie CC, Obare F, RamaRao S, Van Lith L. Communities take action in Kenya: Strengthening postabortion care. New York: EngenderHealth, RESPOND Project; 2013. <http://www.respond-project.org/archive/files/4/4.4/4.4.2/Brief13-2013-Communities-Take.pdf>
- Undie CC, Van Lith LM, Wahome M, Obare F, Oloo E, Curtis C. Community mobilization and service strengthening to increase awareness and use of postabortion care and family planning in Kenya. *International Journal of Gynaecology and Obstetrics*. 2014;126(1):8-13. <https://obgyn.onlinelibrary.wiley.com/doi/10.1016/j.ijgo.2013.12.016>
- USAID. Post Abortion Care-Family Planning. Last updated Feb 16, 216. Accessed Mar 22, 2018. <https://www.usaid.gov/what-we-do/global-health/maternal-and-child-health/post-abortion-care-%E2%80%93-family-planning>
- Wang LF, Puri M, Rocca CH, Blum M, Henderson JT. Service provider perspectives on post-abortion contraception in Nepal. *Culture, Health & Sexuality*. 2016;18(2):221-232. <https://www.tandfonline.com/doi/abs/10.1080/13691058.2015.1073358>
- Wendot S, Scott RH, Nafula I, Theuri I, Ikiugu E, Footman K. Evaluating the impact of a quality management intervention on post-abortion contraceptive uptake in private sector clinics in western Kenya: a pre- and post-intervention study. *Reproductive Health*. 2018;15(1):10. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5775589/pdf/12978_2018_Article_452.pdf
- World Health Organization (WHO). Health worker roles in safe abortion care and post-abortion contraception. Geneva: WHO; 2015. http://www.who.int/reproductivehealth/publications/unsafe_abortion/abortion-task-shifting/en/

- Wildemeersch D, Goldstuck ND. Expulsion and continuation rates after postabortion insertion of framed IUDs versus frameless IUDs - review of the literature. Open Access Journal of Contraception. 2015;6:87-94. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5683144/pdf/oajc-6-087.pdf>
- Xu JS, Dai Y, Jiao N, Qian X, Zhang WH. Systematic review of experiences and effects of integrating post-abortion family planning services into existing health system worldwide. Journal of Reproduction and Contraception. 2015;26(1):31-45. <https://www.popline.org/node/643432>
- Zaidi S, Yasmin H, Hassan L, Khakwani M, Sami S, Abbas T. Replacement of dilation and curettage/evacuation by manual vacuum aspiration and medical abortion, and the introduction of postabortion contraception in Pakistan. International Journal of Gynecology & Obstetrics. 2014;126(S1):S40-S44. <https://obgyn.onlinelibrary.wiley.com/doi/full/10.1016/j.ijgo.2014.03.016#support-information-section>
- Zavier AF, Jejeebhoy SJ. Contraceptive use and intentions among unmarried and married young women undergoing abortion in Bihar and Jharkhand, India. Asia-Pacific Population Journal. 2015;30(1):51-70. <http://www.unescap.org/sites/default/files/APPJ-Vol-30-No-1.pdf>
- Zhang WH, Li J, Che Y, et al. Effects of post-abortion family planning services on preventing unintended pregnancy and repeat abortion (INPAC): A cluster randomised controlled trial in 30 Chinese provinces. Lancet. 2017;390:S29. [http://www.thelancet.com/pdfs/journals/lancet/PIIS0140-6736\(17\)33167-7.pdf](http://www.thelancet.com/pdfs/journals/lancet/PIIS0140-6736(17)33167-7.pdf)
- Zhao W, Li J, Meng Q, Liu J. Integrating post-abortion family planning services into existing abortion services in hospital settings in China (INPAC) project management characteristics. Chinese Journal of Medical Science Research Management. 2016;29(2):104-106. https://cordis.europa.eu/result/rcn/199647_en.html

